

Table 1. Rates of Sexual Intercourse and Failure to Use Condoms Among 383 African American Youth at Time Periods Throughout Three Years Post-intervention

	Baseline		6 Months		12 Months		18 Months		24 Months		36 Months	
	C	I	C	I	C	I	C	I	C	I	C	I
Full Sample												
N (%) Completed the Survey	177 (46)	206 (54)	132 (44)	169 (56)	114 (41)	162 (59)	112 (43)	151 (57)	103 (42)	142 (58)	85 (48)	93 (52)
Had sexual intercourse	69 (40)	67 (34)	35 (27)	46 (29)	44 (39)	49 (30)	45 (41)	54 (36)	43 (42)	61 (44)	37 (46)	43 (49)
Failure to Use Condom ¹	25 (41)	16 (29)	12 (39)	7 (17)*	11 (28)	10 (24)	17 (42)	10 (19)*	12 (29)	20 (36)	10 (29)	9 (25)
Restricted Sample One (N = 305)²												
Had sexual intercourse	48 (36)	60 (35)	32 (24)	53 (31)	50 (38)	51 (30)	54 (41)	64 (37)	59 (45)	79 (46)	63 (48)	83 (48)
Failure to Use Condom ¹	15 (31)	13 (22)	12 (38)	9 (17)*	11 (22)	11 (22)	16 (30)	10 (16)	13 (22)	21 (27)	14 (22)	9 (11)
Restricted Sample Two (N = 229)³												
Had sexual intercourse	27 (29)	41 (31)	16 (17)	34 (26)	30 (32)	33 (25)	37 (40)	49 (37)	38 (41)	55 (42)	40 (43)	61 (46)
Failure to Use Condom ¹	11 (41)	8 (20)	7 (44)	7 (21)	7 (23)	8 (24)	15 (41)	9 (18)*	8 (21)	19 (35)	7 (18)	8 (13)

¹Only assessed among youth who had sexual intercourse at each wave;

²Youth who finished two or more follow-up assessments;

³Youth who finished at least four of the five follow-up assessments.

*P < 0.05; C = control; I = intervention